

Kerb[®] FLO

HERBICIDE

Product Registration Number: MAPP 13716

A suspension concentrate containing 400 g/litre (35.3% w/w) propyzamide.

A residual herbicide for the control of a wide range of weeds in WINTER OILSEED RAPE and several other AGRICULTURAL and HORTICULTURAL crops, and in FORESTRY and AMENITY SITUATIONS.


The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

5 Litres e

**PROTECT FROM FROST
SHAKE WELL BEFORE USE
Dow AgroSciences Limited**

CPC2 Capital Park,
Fulbourn, Cambridge, CB21 5XE
Telephone: 01462 457272 Fax: 01462 426605
24 hour Emergency Telephone Number:
+44 (0) 1553 761 251
For Technical Enquiries Call: 0800 6898899



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]: Kerb[®] Flo, Propyzamide

WARNING

SUSPECTED OF CAUSING CANCER. VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.

Do not handle until all safety precautions have been read and understood.
Wear protective gloves/protective clothing/eye protection/face protection.
IF exposed or concerned: Get medical advice/attention
Collect spillage.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 2-Naphthalenesulfonic acid, 6-hydroxy-, polymer with formaldehyde and methylphenol, sodium salt and 1,2-benzisothiazolin-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/KERB-FLO-400-SDS.pdf>

Alternatively contact your supplier

®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow



Kerb[®] FLO

HERBICIDE

Product Registration Number: MAPP 13716.

A suspension concentrate containing 400 g/litre (35.3% w/w) propyzamide.

A residual herbicide for the control of a wide range of weeds in WINTER OILSEED RAPE and several other AGRICULTURAL and HORTICULTURAL CROPS, and in FORESTRY and AMENITY SITUATIONS.

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying by vehicle-mounted or trailed equipment. WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when applying by hand-held equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS before meals and after work.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal:

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place. EMPTY CONTAINER COMPLETELY and dispose of safely.

PROTECT FROM FROST.

SHAKE WELL BEFORE USE.

BATCH NUMBER:

5 Litres e

This label is compliant with the CPA Voluntary Initiative Guidance



The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/FORESTRY HERBICIDE

Crops/Situations

Oilseed rape (winter), sugar beet (seed crop), field bean (winter), apple, blackberry, blackcurrant, gooseberry, loganberry, pear, plum, raspberry, redcurrant, rhubarb (outdoor use only), strawberry (outdoor use only), red clover (seed crop), white clover (seed crop), fodder rape (seed crop), kale (seed crop), turnip (seed crop), lucerne, lettuce (outdoor use only), forest, farm forestry, forest nursery, hedgerow, ornamental plant production (Christmas trees only), amenity vegetation

Maximum Individual Dose

Maximum Number of Treatments) Full details are given in the Important Information

Latest Time of Application) area on the attached leaflet

Other Specific Restrictions:

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

P990474371709

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

IMPORTANT INFORMATION

FUR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/FORESTRY HERBICIDE

Crops/Situations	Maximum Individual Dose (litres product/hectare)	Maximum Number of Treatments	Latest Time of Application
Oilseed rape (winter), sugar beet (seed crop), field bean (winter)	2.1	One per crop	Before 1 st February in year of harvest
Apple, blackberry, blackcurrant, gooseberry, loganberry, pear, plum, raspberry, redcurrant	4.25	One per year	Before 1 st February in year of harvest
Rhubarb (outdoor use only)	4.25	One per year	Before 1 st January in year of harvest
Strawberry (outdoor use only)	3.5	One per year	Before 1 st January in year of harvest
Red clover (seed crop), white clover (seed crop), fodder rape (seed crop), kale (seed crop), turnip (seed crop), lucerne	1.75	One per crop	Before 1 st February in year of harvest
Lettuce (outdoor use only)	3.5	One per crop	Six weeks before harvest
Forest, farm forestry, forest nursery, hedgerow	3.75	One per year	-
Ornamental plant production (see Other Specific Restrictions)	3.75	One per year	-
Amenity vegetation	4.25	One per year	-

Other Specific Restrictions:

Use in ornamental plant production is restricted to application to Christmas trees only.

This product may only be applied to edible crops except lettuce between 1st October and the specified latest time of application.

DO NOT HARVEST CROPS FOR HUMAN OR ANIMAL CONSUMPTION FOR AT LEAST 6 WEEKS AFTER LAST APPLICATION. READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

WARNINGS

Take care to avoid local overdosing.

Do not make more than one application of KERB FLO within 9 months to the same area of land.

SOIL TYPES

KERB FLO can be used on all soil types with the following exceptions:

1. Do not use on soils containing more than 10% organic matter.
2. Do not use on strawberries or winter field beans grown in certain soil types. See Crop Recommendation tables.

SOIL TEXTURE (85 System)	
Textural Group	Textural class
Sands	Coarse sand, Sand, Fine sand, Loamy coarse sand
Very Light Soils	Loamy sand, Loamy fine sand, Coarse sandy loam
Light Soils	Sandy loam, Fine sandy loam, Sandy silt loam, Silt loam (85)
Medium Soils	Sandy clay loam, Clay loam, Silty clay loam
Heavy Soils	Sandy clay, Clay, Silty clay

SOIL AND WEATHER CONDITIONS

KERB FLO requires moisture for root uptake. Best residual action is obtained in moist soils of fine tilth.

KERB FLO can be applied under frosty conditions but should not be used where run-off from the soil surface is likely.

Best results are achieved when growth of weeds (especially blackgrass and volunteer cereals) is slow, but transpiration continues. In mild autumns/winters, emerged weeds may take longer to be controlled, the residual activity of KERB FLO will be shortened and overall control may be reduced.

The efficacy of KERB FLO may be reduced in organic soils and in the presence of excessive surface organic debris, burnt straw, ash, or ploughed-up turf.

In winter field beans ensure a firm seedbed before spraying.

RESISTANCE

Strains of some annual grasses (eg blackgrass, wild oats, Italian ryegrass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

CROP INFORMATION

Note: Specific application information for each crop is contained in the Crop Recommendation Tables that follow.

WINTER OILSEED RAPE

KERB FLO can be applied after the use of an approved specific graminicide applied in accordance with the manufacturer's recommendations.

CLOVER, BRASSICAS AND SUGAR BEET GROWN FOR SEED

Treated clover, brassicas and sugar beet must not be used for human or animal consumption.

AMENITY VEGETATION – TREES, SHRUBS AND ORNAMENTAL PLANTS

KERB FLO is recommended for use on the following species which have been planted out not later than the previous spring: *Berberis darwinii*, *Buddleia davidii*, *Chamaecyparis spp* (False Cypress), *Cotoneaster spp*, *Crataegus spp* (Hawthorns), *Fagus sylvatica* (Beech), *Forsythia x intermedia*, *Ilex aquifolium* (Holly), *Larix decidua* (Larch), *Philadelphus spp* (Mock Orange), *Picea spp* (Spruces), *Pinus spp* (Pines), *Prunus gladiolosa*, *Quercus spp* (Oak), *Rosa spp* (Roses including Rose rootstocks), *Spiraea x bumalda 'Froebelii'*, *Syringa vulgaris* (Lilac), *Taxus baccata* (Yew), *Thuja orientalis*, 'Rosedalis'.

FORESTRY, FARM FORESTRY, FOREST NURSERY, HEDGEROW

KERB FLO is recommended for use on the following species: Alder, Beech, Southern Beech, Birch, Douglas Fir, Grand Fir, Noble Fir, Horse Chestnut, Larch, Lawson Cypress, Oak, Bishop Pine, Corsican Pine, Lodgepole Pine, Monterey Pine, Scots Pine, Poplar, Norway Spruce, Sycamore, Sitka Spruce, Western Hemlock, Wild Cherry.

ORNAMENTAL PLANT PRODUCTION

KERB FLO may be used in the production of Christmas trees only.

PROCESSED CROPS

No taints have been detected in tests with treated field crops. Consult your processor before use.

WATER VOLUMES

Winter oilseed rape, sugar beet (seed crops), winter field beans, lucerne, brassicas (for seed production), clover (seed crops): Apply KERB FLO in 200 to 1000 litres of water per hectare.

Apple, blackcurrant, blackberry, gooseberry, loganberry, pear, plum, raspberry, redcurrant, rhubarb (outdoor), strawberry (outdoor), lettuce (outdoor), forestry, farm forestry, forest nursery, hedgerow, ornamental plant production (Christmas trees only), amenity vegetation: Apply KERB FLO in 400 to 1000 litres of water per hectare.

Ensure good ground cover.

APPLICATION EQUIPMENT

Do not apply through broadcast air-assisted sprayers.

All crops: Application may be made through a ground crop sprayer.

Forestry, farm forestry, forest nursery, hedgerow, ornamental plant production (Christmas trees only), amenity vegetation: KERB FLO may also be applied via a knapsack sprayer.

FOLLOWING CROPS

The number of weeks which must elapse between the last application of KERB FLO and drilling or planting of the following crop is shown in the table below:

Following crops	Rate of KERB FLO applied to previous crop	Date KERB FLO applied to previous crop	
		1 st April – 31 st July	1 st August – 31 st March
Lettuce	1.75–4.25 litres/ha	0 weeks	0 weeks
Field beans, broad beans, peas, chicory, radish, clover, lucerne	1.75–4.25 litres/ha	5 weeks	10 weeks
Brassicas, leeks, onions, parsley, parsnips, celery, oilseed rape, strawberries	1.75–4.25 litres/ha	10 weeks	25 weeks or plant/sow after 15 th June whichever occurs sooner
Cereals and grasses ⁴	1.75–2.1 litres/ha	40 weeks	30 weeks
	2.75–4.25 litres/ha		40 weeks
Any other crop ⁴	2.75–4.25 litres/ha	20 weeks	40 weeks

⁴Treated land must be mouldboard ploughed to a depth of 15 cm prior to drilling a following cereal, grass or any other crop not listed above.

Please consult Dow AgroSciences if a treated crop fails because of bad growing conditions.

MIXING

Add half the required volume of water to the spray tank and begin agitation. Shake the container vigorously and add the recommended quantity of KERB FLO through the filter basket. When container becomes empty, wash out with water and add the washings through the filter basket. Rinse through the filter basket and add the rest of the water. Agitate while topping up the tank and continue agitation until spraying is complete.

Spray immediately: do not allow the mixture to stand.

Thoroughly wash all spraying and measuring equipment with water immediately after use.

CROP RECOMMENDATION TABLES

S = Susceptible MS = Moderately susceptible MR = Moderately resistant R = Resistant

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germi-nating	Up to 2 leaf	Estab-lished			
Winter oilseed rape	1.75 litres/ha	Annual meadow-grass, barren brome, volunteer cereals, wild-oat	S	S	S	1 st October to 31 st January	As soon as possible after 3rd true leaf stage. Crop selectivity is by depth protection. Factors which cause shallow rooting may reduce crop selectivity.	All soils with less than 10% organic matter.
		Common chickweed	S	S	S ¹			
		Blackgrass	S	S	MS ²			
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MR			
		Field forget-me-not	MS	MS	R			
	Cleavers	MS	MR	R				
	1.25 litres/ha - Use this rate only where a specific graminicide has controlled volunteer cereals and grass weeds and chickweed is not a problem	Annual meadow-grass, volunteer cereals, wild-oat	S	S	S			
Black-bindweed, black nightshade, common chickweed, fat-hen, knotgrass, redshank, small nettle		S	S	R				
Speedwells		S	R	R				
Sugar beet grown for seed	2.1 litres/ha	Annual meadow- grass, barren brome, common chickweed, volunteer cereals, wild-oat	S	S	S	Sugar beet grown for seed: 1 st October to 31 st January	Sugar beet grown for seed: As soon as possible after 4th true leaf stage.	Sugar beet grown for seed: All soils with less than 10% organic matter.
		Blackgrass	S	S	MS			
Winter field beans		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MR	Winter field beans: 1 st October to 31 st January	Winter field beans: Within 7 days after drilling but before crop emerges.	Winter field beans: Use only on medium or heavy soils with less than 10% organic matter.
		Cleavers	MS	MR	R			

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germinating	Up to 2 leaf	Established			
Apple, blackcurrant, blackberry, gooseberry, loganberry, pear, plum, raspberry, redcurrant, rhubarb (outdoor use only), strawberry (outdoor use only)	2.1 litres/ha	Blackgrass	S	S	MS	Strawberry, rhubarb: 1 st October to 31 st December Other crops: 1 st October to 31 st January	Strawberry: Established crops planted for at least one year. Other crops: Established crops planted for at least one season.	Strawberry: Use only on heavy soils with less than 10% organic matter. Other crops: All soils with less than 10% organic matter.
		Annual meadow-grass, barren brome, common chickweed, volunteer cereals, wild-oat	S	S	S			
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MR			
	Strawberry: (not matted row crops) 3.5 litres/ha Other crops: 4.25 litres/ha	Annual meadow-grass, barren brome, blackgrass, common chickweed, volunteer cereals, wild-oat	S	S	S			
		Common couch ³ and other perennial grasses	S	S	S			
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MS			
		Cleavers	S	S	R			
		Common fumitory, shepherd's-purse	MS	MS	R			
		Creeping buttercup, broad-leaved dock, sheep's sorrel	S	MS	MS			
		Field horsetail	MS	MS	MS			

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germi-nating	Up to 2 leaf	Estab-lished			
Brassicas grown for seed (fodder rape, kale, turnip), clover grown for seed (red and white), lucerne	1.75 litres/ha	Blackgrass	S	S	MS	1 st October to 31 st January	<p>Brassicas grown for seed: As soon as possible after 3rd true leaf stage.</p> <p>Clover grown for seed: Established crops grown for at least one season.</p> <p>Lucerne: Established crops grown for at least one season. Not less than 7 days after last cut.</p>	All soils with less than 10% organic matter.
		Annual meadow-grass, barren brome, common chickweed, volunteer cereals, wild-oat	S	S	S			
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MR			
		Cleavers	MS	MR	R			
Lettuce (outdoor use only)	2.75-3.5 litres/ha Under dry soil conditions or warm weather use 3.5 litres/ha and irrigate or incorporate.	Annual meadow-grass, barren brome, blackgrass, common chickweed, volunteer cereals, wild-oat	S	S	S	Any time up to 6 weeks before harvest	Before or after drilling	All soils with less than 10% organic matter.
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MS			
		Cleavers, common fumitory, shepherd's-purse	MS	MS	R			

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germinating	Up to 2 leaf	Established			
Forestry, farm forestry, forest nursery, hedgerow (see list of species under 'Crop Information') Ornamental plant production (Christmas trees only)	3.75 litres/ha	Common bent, common couch, sweet vernal grass, tufted hair-grass, Yorkshire-fog and other perennial grasses	S	S	S	1 st October to 31 st January North of a line Aberystwyth to London (see also Soil Type)	Forests: Any stages including pre-planting. Nurseries: Treat no earlier than one month after transplanting.	Mineral Soils (surface water gleys, brown earths and sands). Use between 1 st October and 31 st January. (See also Time of Year). Peat Soils (peaty gleys and peat soils with a depth of organic matter greater than 10 cm): between 1 st October and 31 st December. (See also Time of Year).
		Creeping soft-grass	S	S	MS			
		Cock's foot	S	S	MR			
		Field horsetail	MS	MS	MS			
		Sedges	MS	MS	MS			
		Foxglove, willowherbs	R	R	R	1 st October to 31 st December South of a line Aberystwyth to London (see also Soil Type)		

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germinating	Up to 2 leaf	Established			
Amenity vegetation – trees, shrubs and ornamental plants (see list of species under 'Crop Information')	2.1 litres/ha	Annual meadow-grass, barren brome, blackgrass, common chickweed, volunteer cereals, wild-oat	S	S	S	1 st October to 31 st January	Established crops planted for at least one season.	All soils with less than 10% organic matter.
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MR			
	4.25 litres/ha	Annual meadow-grass, barren brome, blackgrass, common chickweed, volunteer cereals, wild-oat	S	S	S			
		Common couch and other perennial grasses	S	S	S			
		Black-bindweed, black nightshade, fat-hen, knotgrass, redshank, small nettle, speedwells	S	S	MS			
		Cleavers	S	S	R			
		Common fumitory, shepherd's-purse	MS	MS	R			
		Creeping buttercup, broad-leaved dock, sheep's sorrel	S	MS	MS			
		Field horsetail	MS	MS	MS			
All listed crops	All rates	Common poppy, gallant soldier, groundsel, mayweed, scarlet pimpernel, clover, dandelion, field bindweed, ragwort, thistle	R	R	R	Any time	Any stage	All soils

NOTES FOR CROP RECOMMENDATION TABLES

1	Chickweed control may be reduced where it is well established (over 10 cm in diameter).
2	CONTROL OF BLACKGRASS IN WINTER OILSEED RAPE Established (well-tillered) blackgrass is moderately susceptible: Where populations of blackgrass and/or volunteer cereals exceed 50/m ² KERB FLO should be applied in tank mix with an approved graminicide, or following an effective approved graminicide to ensure optimum weed control. Deeper germinating blackgrass within the soil profile could reduce product efficacy. When applied alone in late season, it is unlikely this higher dose will markedly improve control of well tillered blackgrass. Where partial resistance (R [†] or RR [†]) to the partner graminicide is known to exist the dose of KERB FLO may be increased to 2.1 litres/ha. This may also be done if applications are made early in the season, under warm conditions and an increase in the duration of residual control is required (see 'Soil and weather conditions' above). Where high levels of resistance (RRR [‡]) to the partner graminicide occurs there is no advantage of adding this graminicide to KERB FLO for blackgrass control, and KERB FLO even at 2.1 litres/ha will not give acceptable levels of established blackgrass control in these circumstances. [†] R = 1* RR = 2*/3* RRR = 4*/5*
3	For heavy infestation of common couch a repeat application may be required the following winter. Avoid deep burying of rhizomes when preparing land prior to KERB FLO treatment.

TRADEMARK ACKNOWLEDGEMENTS

KERB is a trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow.

All other brand names used in this document are trademarks of other manufacturers in which proprietary rights may exist.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN



Dow AgroSciences



Kerb® FLO

HERBICIDE

Product Registration Number: MAPP 13716.

A suspension concentrate containing 400 g/litre (35.3% w/w) propyzamide.

A residual herbicide for the control of a wide range of weeds in WINTER OILSEED RAPE and several other AGRICULTURAL and HORTICULTURAL CROPS, and in FORESTRY and AMENITY SITUATIONS.

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying by vehicle-mounted or trailed equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND RUBBER BOOTS when applying by hand-held equipment.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS before meals and after work.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal:

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

PROTECT FROM FROST.

SHAKE WELL BEFORE USE.

BATCH NUMBER:

5 Litres e

This label is compliant with the CPA Voluntary Initiative Guidance



The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/FORESTRY HERBICIDE

Crops/Situations

Oilseed rape (winter), sugar beet (seed crop), field bean (winter), apple, blackberry, blackcurrant, gooseberry, loganberry, pear, plum, raspberry, redcurrant, rhubarb (outdoor use only), strawberry (outdoor use only), red clover (seed crop), white clover (seed crop), fodder rape (seed crop), kale (seed crop), turnip (seed crop), lucerne, lettuce (outdoor use only), forest, farm forestry, forest nursery, hedgerow, ornamental plant production (Christmas trees only), amenity vegetation

Maximum Individual Dose

Maximum Number of) Full details are given in the Treatments Important Information

Latest Time of Application) area on the attached leaflet

Other Specific Restrictions:

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow